

Teething

QUALITY AREA 2: CHILDREN'S HEALTH AND SAFETY

Policy Statement

Our Service aims to manage the teething process of babies and young children whilst ensuring the health and safety of all children, families, educators and visitors of the Service. Symptoms of common childhood illnesses are often mistakenly linked to 'teething', which may prevent the identification of an underlying bacterial or viral or middle ear infection.

This policy aims to ensure staff, parents, families and visitors to the Service understand teething symptoms, measures implemented to manage the teething process and treatments that are not recommended or used at our Service.

Aims

We aim to assist families manage the teething process of their child whilst maintaining the health and wellbeing of all children.

Strategies

Our Service implements specific strategies to ensure the health and wellbeing of all children, families, educators and visitors. Our policies are informed by current research and recommendations provided by the Australian Government-Department of Health and local Public Health Units and we adhere to the recommendations by the Australian Government National Health and Medical Research Council Publication *Staying healthy: Preventing infectious diseases in early childhood education and care services* (Fifth Edition). Although some babies and young children may display signs of discomfort and pain whilst teething, many babies experience little or no discomfort. Our educators will implement a range of management strategies to support and comfort babies and young children during this time.

Teething

Baby teeth develop while babies are still in the womb. Teething is the process in which the teeth begin to 'erupt' and break through the gums. Teething commonly begins between four and ten months of age and is different for each child. The order of tooth eruption however is usually the same with the two front teeth (central incisors) in the lower jaw occurring between the ages of six and ten months.

Generally, children will have their full set of primary teeth (20 teeth) by the age of three years.

Symptoms and/ or behaviours of teething

Teething takes about eight days for a tooth to fully erupt from the gum. During this time babies and young children may suffer some discomfort. Symptoms and/or behaviours may include:

- babies rubbing their gums together

- flushed red cheeks
- sucking on toys, fingers or fists
- irritable, grizzly or grumpy
- baby being more clingy or fretful than usual
- dribbling as more saliva is produced during teething
- nappy rash

Whilst teething symptoms for some babies can be concerning, there is no clear evidence to suggest that teething causes fever or diarrhoea in children (Royal Children's Hospital Melbourne). Symptoms of common childhood illnesses can sometimes be mistakenly linked to teething resulting in underlying viral, bacterial or middle ear infections to go undiagnosed. Teething may cause discomfort and irritability, but it does not cause an illness. If a baby or child has a temperature at or above 38°C parents should seek urgent medical attention.

Management of teething

Educators will implement a range of techniques to comfort and support babies and young children during teething. These may include:

- use a cold, wet flannel (face washer) to massage the gum area with a finger
- if baby has started solid foods and is over six months, offering food items that are firm- such as a sugar free rusk, or raw vegetables
- offering mushier foods for mealtimes (mashed vegetables, yoghurt)
- additional comforting and play time to distract the baby from any pain

Medication for teething

As per our *Incident, Injury, Trauma and Illness Policy*, management will not accept a child into care if they have been given medication for a pain relief or temperature prior to arriving at the Service (for example Panadol). If a child becomes ill whilst at the Service and has developed a temperature, we will adhere to our *Incident, Injury, Trauma and Illness Policy* and monitor the child's symptoms, temperature and respond accordingly.

If the child is displaying signs of pain and discomfort and other measures have not helped, we will administer paracetamol only with written authorisation by a parent. Any medication administered to a child will be recorded on the *Administration of Medication Record* or *Administration of Paracetamol Record* and witnessed by another educator. Dosage amounts and instructions will be checked carefully and recorded appropriately. The *Administration of Medication Record* will be acknowledged and signed by the parent or authorised nominee when collecting the child from the Service. When paracetamol has been administered, it is our policy to request parents or an authorised person to collect the child from care within 30 minutes.

Teething Gels- Our Service will NOT administer teething gels for a child who is teething. Common teething gel formulations contain choline salicylate which is related to aspirin and is not recommended for children under 16 years of age. Teething gels containing benzocaine are also not recommended. (Victoria Health-better health).

Amber Beads- The use of amber beads to assist for soothing teething are not supported by our Service. The Australian Competition and Consumer Complaints (ACCC) has identified amber beads as public safety warning due to the risk of amber teething necklaces and bracelets becoming a possible choking or strangulation hazard.

The Approved Provider, Nominated Supervisor, and Educators will ensure:

- they have knowledge of and adhere to this policy and associated procedure
- they aware of a range of strategies to support babies and young children during teething
- communication with parents and families is consistent and supportive
- children will not be accepted into care if they have been provided with paracetamol or other medication for pain relief or temperature prior to arriving at the Service
- families are advised that amber teething necklaces and bracelets are not to be worn by babies and young children at our Service- as per warnings by the Australian Government (2011) and Red Nose Australia (2025)
- effective hygiene policies and procedures are adhered to at all times to prevent the spread of illnesses
- effective environmental cleaning policies and procedures are adhered to all times
- toys and other equipment mouthed by babies or young children are thoroughly cleaned after use
- all families are provided access to relevant policies upon enrolment which will be explained by management including: *Teething Policy, Incident, Injury, Trauma and Illness Policy, Administration of Medication Policy*
- families are provided with relevant information from a trusted source about teething and management of teething
- that any child who registers a temperature of 38°C or above is to be collected from the Service within 30 minutes
[Failure to comply to this time restriction may result in further action taken by management and regarded as a breach of service policy]
- as per our *Incident, Injury, Trauma and Illness Policy*, children are not to return to the Service if they still have a temperature above 38°C the following day
- administration of medication for pain relief will only occur after all other measures to reduce pain and discomfort for teething children have been exhausted and with written authorisation by a parent or authorised nominee
- if paracetamol is administered to a child for any reason, parents or an authorised person must collect the child within 30 minutes from the Service
- accurate records for *Administration of Medication* and/or *Administration of Paracetamol* are completed, witnessed and signed by a parent/authorised nominee when collecting the child.

Families will:

- adhere to the *Teething Policy and Incident, Injury, Trauma and Illness Policy*
- be informed that amber teething bracelets or necklaces are not to be worn at our Service due to the risk of choking
- not administer paracetamol or other medication for pain relief or temperature prior to arriving to the Service
- provide written authorisation to administer pain relief medication (e.g., Panadol)
- acknowledge and sign the *Administration of Medication Record* if required
- collect their child from the Service within 30 minutes if paracetamol has been administered
- monitor their child's health and recognise the symptoms of an underlying illness rather than teething

Continuous Improvement/Reflection

The Teething policy will be evaluated and reviewed on an annual basis or earlier if there are changes to legislation, ACECQA guidelines or any incident related to our policy. Feedback will be requested from families, staff, educators and management, and notification of any change to policies will be made to families within 14 days.

NATIONAL QUALITY STANDARD (NQS)

QUALITY AREA 2: CHILDREN'S HEALTH AND SAFETY

2.1	Health	Each child's health and physical activity is supported and promoted
2.1.1	Wellbeing and comfort	Each child's wellbeing and comfort is provided for, including appropriate opportunities to meet each child's needs for sleep, rest and relaxation.
2.1.2	Health Practices and Procedures	Effective illness and injury management and hygiene practices are promoted and implemented

EDUCATION AND CARE SERVICES NATIONAL REGULATIONS

77	Health, hygiene and safe food practices
85	Incident, injury, trauma and illness policies and procedures
87	Incident, injury, trauma and illness record
92	Medication record
93	Administration of medication

Source:

Australian Dental Association Oral tips for young children.

Australian Children's Education & Care Quality Authority. (2023). *Guide to the National Quality Framework*.

Education and Care Services National Law Act 2010. (Amended 2023).

Education and Care Services National Regulations. (Amended 2023).

National Health and Medical Research Council. (2013). *Staying Healthy: Preventing infectious diseases in early childhood education and care services* (5th Ed.). Australia: Commonwealth of Australia. NSW Government. (n.d.).

Pregnancy, Birth and Baby <https://www.pregnancybirthbaby.org.au/teething>

Raising Children Network: <https://raisingchildren.net.au/babies/health-daily-care/dental-care/dental-care-babies>

Teeth.org.au Babies and toddlers 0-3 years

The Royal Children's Hospital Melbourne. https://www.rch.org.au/kidsinfo/fact_sheets/Fever_in_children/ Victoria Health Department.

Better Health Channel. Teeth Development in Children.

<https://www.betterhealth.vic.gov.au/health/conditionsandtreatments/teeth-development-in-children> Western

Australian Education and Care Services National Regulations

Review

Policy reviewed by: Suzi Scott

Policy reviewed: July 2026

Next review date; July 2027

Modifications: Update on continuous improvement and reflection, update on information regarding Red Nose Australia